



International Clostridium difficile Animal Strain Collection

Zidaric, V.; Janezic, S.; Pardon, B.; Indra, A.; Kokotovic, Branko; Blanco, J. L.; Seyboldt, C.; Diaz, C. Rodriguez ; Poxton, I. R.; Perreten, V.

Total number of authors:

16

Publication date:

2012

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Zidaric, V., Janezic, S., Pardon, B., Indra, A., Kokotovic, B., Blanco, J. L., Seyboldt, C., Diaz, C. R., Poxton, I. R., Perreten, V., Drigo, I., Jiraskova, A., Ocepek, M., Weese, J. S., Songer, J. G., & Rupnik, M. (2012). *International Clostridium difficile Animal Strain Collection*. Poster session presented at 4th International Clostridium difficile Symposium (ICDS 2012), Bled, Slovenia.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

INTERNATIONAL *Clostridium difficile* ANIMAL STRAIN COLLECTION

V. Zidaric, S. Janezic¹, B. Pardon², A. Indra³, B. Kokotovic⁴, J.L. Blanco⁵, C. Seyboldt⁶, C. Rodriguez Diaz⁷, I.R. Poxton⁸, V. Perreten⁹, I. Drigo¹⁰, A. Jiraskova¹¹, M. Ocepek¹², J.S. Weese¹³, J.G. Songer¹⁴, M. Rupnik^{1, 15,16}

¹Institute for Public Health Maribor, Maribor, Slovenia; ²Ghent University, Faculty of Veterinary Medicine, Merelbeke, Belgium; ³AGES, Vienna, Austria; ⁴Technical University of Denmark, National Veterinary Institute, Copenhagen, Denmark; ⁵Complutense University, Madrid, Spain; ⁶Friedrich-Loeffler Institute, Jena ,Germany; ⁷University of Liege, Faculty of Veterinary Medicine, Liege, Belgium; ⁸University of Edinburgh, Edinburgh, UK; ⁹University of Bern, Institute of Veterinary Bacteriology, Bern, Switzerland; ¹⁰IZSVe, Treviso, Italy; ¹¹Charles University in Prague, 1st Faculty of Medicine, Prague, Czech Republic; ¹²University of Ljubljana, Veterinary Faculty, Ljubljana, Slovenia; ¹³University of Guelph, Ontario Veterinary College, Ontario, Canada; ¹⁴Iowa State University, Ames, U.S.A.; ¹⁵University of Maribor, Medical Faculty, Maribor, Slovenia; ¹⁶Centre of excellence for integrated approaches in chemistry and biology of proteins, Ljubljana, Slovenia

Background

Animals have been recognized as an important potential reservoir of *Clostridium difficile* and according to recent studies the overlap between PCR ribotypes of human and animal isolates seems to be increasing (Bakker *et al.*, 2010; Gould and Limbago, 2010; Janezic *et al.*, 2012; Keel *et al.*, 2007; Koene *et al.*, 2011).

Here we report on an International *Clostridium difficile* animal strain collection that was established to enhance comparative studies on animal-associated strains and contribute to interlaboratory exchange of the strains.

The goal of the collection is to include one PCR ribotype per species per country/laboratory.

Results and Discussion

All 100 included strains were distributed into 39 different PCR ribotypes. Up to 17 different PCR ribotypes can be found within a single animal species and up to 15 different PCR ribotypes per country.

Within one standard (agarose based) PCR ribotype several PCR ribotypes could be distinguished by capillary gel based PCR ribotyping (Table 1; PCR ribotypes 014/020, 078, 002, 045, 056, 033).

Five strains are nontoxigenic while toxigenic strains account for 95.0% and belong to 10 different toxinotypes: 0, I, III, IV, V, VI, VIII, XI, XII and XIX. PCR ribotypes 078, 126, 014/020, 012 and 002 that are frequently associated with animals (Keel *et al.*, 2007; Janezic *et al.*, 2012) represent 40.0 % of all strains.

Materials and methods

C. difficile strains:

Altogether 100 strains from 12 different countries were contributed. Collected strains originate from 11 different animal species, including pets, horses and food animals. Approximately half (58,0%) of the strains are from cattle and pigs (Table 1).

For every strain additional available information on animal host and strain was obtained.

Molecular characterization of *C. difficile* strains:

All collected strains were characterized by toxinotyping (Rupnik *et al.*, 1998; <http://www.mf.uni-mb.si/tox/>). In addition, binary toxin genes were detected by PCR as described in Stubbs *et al.* (2000).

Standard agarose gel-based PCR ribotyping was used as described by Bidet *et al.* (1999) and results analyzed by BioNumerics software 5.10 (Applied Maths).

Strains were also typed by capillary gel electrophoresis-based ribotyping using primers for standard agarose gel-based PCR ribotyping with flourescein labelled 16S primer (Indra *et al.*, 2008). PCR ribotype patterns were analyzed and identified on a web-based database Webribo (<http://webribo.ages.at>).

Table 1. Distribution of *C. difficile* genotypes by species and country given as toxinotype/standard PCR ribotype [capillary gel electrophoresis based PCR ribotype].

Country	Austria	Belgium	Canada	Czech Republic	Denmark	Germany	Italy	Scotland	Slovenia	Spain	Switzerland	U.S.A.	Total number of strains per animal species	Number of different PCR ribotypes per animal species
Animal host														
Pig	XII/056 [new RT] 0/011/049 [049/1] 0/150 [AI-12]	V/078 [078] 0/002 [002/0] 0/081 [081]	V/078 [078] 0/015 [211]	0/150 [AI-12]	V/078 [new RT] V/045 [598] V/126 [126] VI/066 [413] 0/150 [AI-12] 0/005 [005] VI/SLO 137 [new RT]		V/078 [251] 0/150 [AI-12] XIX/012 [012] 0/014/020 [014 subtype] 0/081 [081] 0/SLO 036 [050/AI-84] XII/SLO 133 [AI-15] VI/SLO 012 [new RT]	V/078 [078]	V/045 [045]	V/078 [078]		V/078 [078] 0/002 [nd]	29	17
Cattle	V/078 [078] 0/014/020 [014/0] 0/029 [029] 0/005 [005] 0/015 [211] 0/SLO 125 [014 subtype] 0/SLO 036 [AI-84/050] 0/SLO 090 [AI-9-1] 0/SLO 143 [610] 0/SLO 164 [nd] 0/SLO 165 [nd]	V/078 [078] V/078 [078] V/126 [078 ecdc] XI/033 [033] 0/012 [012] 0/014/020 [014/0] 0/002 [002/2] 0/081 [081]	0/SLO 074 [new RT] V/078 [078]				V/126 [126] XIIb/033 [033]		0/002 [209] 0/014/020 [014/0]	XIIb/283 [new RT]		V/078 [078] XIa/033 [033] III/027 [027]	29	15
Dog			0/001/072 [nd] 0/002 [002/0]			0/SLO 066 [AI-60] 0/014/020 [014/0] V/SLO 024 [new RT] Tox-/010 [010]			0/014/020 [new RT] 0/012 [012]	XII/056 [new RT] Tox-/010 [010]			10	8
Cat			0/001/072 [001 ecdc]			V/SLO 024 [new RT] 0/014/020 [449] Tox-/SLO 002 [AI-34]			0/014/020 [new RT]				5	4
Poultry	0/001/072 [001]								0/029 [029] 0/014/020 [014/0] IV/023 [023] 0/103 [AI-82/1] 0/001/072 [001] XI/033 [nd]				6	5
Horse		0/014/020 [020]	XIIb/033 [new RT] V/078 [078] III/027 [027]								V/078 [078] V/126 [126]		7	5
Rabbit/Hare							0/012 [012] V/078 [078] 0/014/020 [014/0] Tox-/SLO 132 [new RT] 0/SLO 090 [AI-9-1] 0/002 [002/0] VIII/017 [017] Tox-/SLO 084 [205]						8	8
Partridge									XII/056 [446] XI/033 [new RT]				2	2
Goose									0/SLO 090 [AI-9-1]				1	1
Crow									0/003 [003]				1	1
Raccoon			I/SLO 166 [nd] 0/103 [AI-82/1]										2	2
Total number of strains per country	15	12	12	1	7	7	18	1	16	4	2	5	100	
Number of different PCR ribotypes per country	15	7	9	1	7	5	15	1	12	4	2	4		

nd- typing not done ; new RT- PCR ribotype has on webribo database not yet been determined; subtype- subtype has on webribo not yet been determined

References

Bakker D, Corver J, Harmanus C, Goorhuis A, Keessen EC, Fawley WN, Wilcox MH, Kuijper EJ. 2010. Relatedness of human and animal *Clostridium difficile* PCR ribotype 078 isolates determined on the basis of Multilocus Variable-Number Tandem-Repeat Analysis and tetracycline resistance. *J. Clin. Microbiol.* **48**(10):3744–3749.

Bidet P, Barbut F, Lalande V, Burghoffer B, Petit JC. 1999. Development of a new PCR-ribotyping method for *Clostridium difficile* based on ribosomal RNA gene sequencing. *FEMS Microbiol. Lett.* **175**:261–266.

Gould LH, Limbago B. 2010. *Clostridium difficile* in food and domestic animals: a new foodborne pathogen? *Clin Infect Dis.* **51**(5):577–582.

Indra A, Huhulescu S, Schneeweis M, Hasenberger P, Kerubichler S, Fiedler A, Wewalka G, Allerberger F, Kuijper EJ. 2008. Characterization of *Clostridium difficile* isolates using capillary gel electrophoresis-based PCR ribotyping. *J. Med. Microbiol.* **57**: 1377–1382.

Janezic S, Ocepek M, Zidaric V, Rupnik M. 2012. *Clostridium difficile* genotypes other than ribotype 078 that are prevalent among human, animal and environmental isolates. *BMC Microbiol.* **12**:48 doi:10.1186/1471-2180-12-48.

Keel K, Brazier JS, Post KW, Weese S, Songer JG. 2007. Prevalence of PCR ribotypes among *Clostridium difficile* isolates from pigs, calves, and other species. *J Clin. Microbiol.* **45**(6):1963–1964.

Koene MG, Mevius D, Wagenaar JA, Harmanus C, Hensgens MP, Meetsma AM, Putirulan FF, van Bergen MA, Kuijper EJ. 2011. *Clostridium difficile* in Dutch animals: their presence, characteristics and similarities with human isolates. *Clin. Microbiol. Infect.* doi:10.1111/j.1469-0691.2011.03651.x

Rupnik M, Avesani V, Janc M, Von Eichel-Streiber C, Delmée M. 1998. A novel toxinotyping scheme and correlation of toxinotypes with serogroups of *Clostridium difficile* isolates. *J. Clin. Microbiol.* **36**(8):2240–2247.

Stubbs S, Rupnik M, Gilbert M, Brazier J, Duerden B, Popoff M. 2000. Production of actin-specific ADP-ribosyltransferase (binary toxin) by strains of *Clostridium difficile*. *FEMS Microbiol. Lett.* **186**:307–312.